

Hiroyoshi OHASHI*: New names for two Asiatic legumes

大橋広好*: アジア産マメ科植物の2新学名

During my studies on Himalayan legumes in the 1970's I proposed two new combinations for the names of varieties of *Bauhinia scandens* (Caesalpinoideae-Cercideae) and *Shutteria involucrata* (Papilionoideae-Phaseoleae). They are *B. scandens* var. *horsfieldii* (Miq.) Ohashi and *S. involucrata* var. *glabrata* (Wight & Arn.) Ohashi. These two varieties were named on the basis of *Bauhinia anguina* var. *horsfieldii* (Miq.) Prain and *Shutteria vestita* var. *glabrata* (Wight & Arn.) Baker, respectively. However, both names must be replaced by the autonym due to the article 57.3 of the International Code of Botanical Nomenclature which is added since 1983. According to this article the autonym of *B. anguina* var. *horsfieldii*, i.e., var. *angula*, has priority over the previous variety name, and that of *S. vestita* var. *glabrata*, i.e., var. *vestita*, is treated as having priority over var. *glabrata*.

The nomenclatural treatments are as follows:

1) ***Bauhinia scandens* L. var. *anguina* (Roxb.) Ohashi, comb. nov.**

Bauhinia anguina Roxb., [Hort. Bengal. 31 (1814), n.n.] Pl. Corom. 3: 82, t. 285 (1819).

Lasiobema horsfieldii Miq., Fl. Ind. Bat. 1(1): 71 (1855).

B. anguina var. *horsfieldii* (Miq.) Watt. ex Prain in J. Asiat. Soc. Beng. 66-2: 194 (1897).

B. horsfieldii (Miq.) Macbride in Contr. Gray Herb. 2, no. 59: 23 (1919).

B. subrhombicarpa Merr. in Lingn. Sci. J. 14: 9 (1935).

L. scandens (L.) De Wit var. *horsfieldii* (Prain) De Wit in Rainwardtia 3: 437 (1956).

B. scandens var. *horsfieldii* (Miq.) K. Larsen & S.S. Larsen in Nat. Hist. Bull. Siam Soc. 25: 16 (1973); Fl. Camb., Laos & Vietn. 18: 168 (1980). Ohashi in Hara & Williams, Enum. Flow. Pl. Nepal 2: 108 (1979), ut var. *horsfieldii* (Miq.) Ohashi.

Distr. Tropical E. Himalaya (Nepal to Bhutan), India, Indo-China, Malaysia.

* Biological Institute, Faculty of Science, Tohoku University, Sendai 980. 東北大学 理学部生物学教室.

2) *Shutteria involucrata* (Wall.) Wight & Arn. var. *vestita* (Wight & Arn.) Ohashi, comb. nov.

[*Glycine vestita* Grah. in Wall., Cat. 190, n. 5512 (1831-32), n.n.]

S. vestita Wight & Arn., Prodr. Fl. Ind. Or. 1: t. 165 (1834). Baker in Hook. f., Fl. Brit. Ind. 2: 181 (1876). Prain in J. Asiat. Soc. Beng. 66-2: 401 (1897). Merr. in Philip. J. Sci. Bot. 5: 110 & 354 (1910); Enum. Philip. Pl. 304 (1923). Gagnep. in Fl. Gén. Indoc. 2: 403 (1916). Craib, Fl. Siam. Enum. 1: 438 (1928). Back. & Bakh. f., Fl. Java 1: 625 (1963). Ohashi in Fl. East. Himal. 163 (1966) & 3: 69 (1975). Icon. Cormophyt. Sin. 2: 492 (1972). Van Thuan in Adansonia ser. 2, 12: 295 (1972); in Fl. Camb. Laos Vietn. 17: 64 (1979). Verdcourt, Man. New Guinea Leg. 480 (1979), in nota.

S. glabrata Wight & Arn., Prodr. Fl. Ind. Or. 207 (1834).

S. vestita var. *glabrata* (Wight & Arn.) Baker in Hook. f., Fl. Brit. Ind. 2: 182 (1876).

S. involucrata var. *glabrata* (Wight & Arn.) Ohashi in Journ. Jap. Bot. 50: 305 (1975); in Hara & Williams, Enum. Flow. Pl. Nepal 2: 130 (1979). Index Fl. Yunnan. 1: 628 (1984). Fl. Xizangica 2: 742 (1985).

f. *vestita* (Wight & Arn.) Ohashi, stat. nov.

f. *densiflora* (Benth.) Ohashi in Journ. Jap. Bot. 50: 305 (1975).

S. densiflora Benth. in Miq., Pl. Jungh. 232 (1852). Prain, l.c. 402 (1897).

S. vestita var. *densiflora* (Benth.) Baker, l.c. 182 (1876). Ohashi, l.c. 3: 69 (1975).

Distr. Himalaya (Simla to Bhutan), India, Ceylon, S.E. Asia, Malaysia, China.

* * * *

国際植物命名規約 57.3 条に基づいて、新学名 *Bauhinia scandens* L. var. *anguina* (Roxb.) Ohashi および *Shutteria involucrata* (Wall.) Wight & Arn. var. *vestita* (Wight & Arn.) Ohashi を発表した。両変種は東南アジアに広く分布する。